

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Tzu Chi Medical Journal

journal homepage: www.tzuchimedjnl.com

Case Report

Mixed fungal keratitis of *Penicillium* species and *Acremonium* speciesNancy Chen^a, Yuan-Chieh Lee^{a,b,c,*}^a Department of Ophthalmology, Buddhist Tzu Chi General Hospital, Hualien, Taiwan^b Department of Medicine, Tzu Chi University, Hualien, Taiwan^c Department of Ophthalmology, National Taiwan University Hospital, Taipei, Taiwan

ARTICLE INFO

Article history:

Received 22 March 2010

Received in revised form

27 April 2010

Accepted 6 May 2010

Keywords:

Acremonium species

Mixed fungal keratitis

Penicillium species

ABSTRACT

Mixed fungal keratitis of two different genera has rarely been reported. We described a case of mixed fungal keratitis of *Penicillium* sp and *Acremonium* sp. A 67-year-old male developed severe keratitis after undergoing superficial corneal trauma of his right eye while picking herbs (*Ficus erecta* Thunb. var.) 3 days earlier. Corneal stromal infiltration with a feathery border and a surrounding immunological reaction ring was noted without epithelial defect. Direct smear examination of corneal scraping showed the presence of filamentous fungi, and topical natamycin (5%) hourly was prescribed along with systemic voriconazole. However, hypopyon soon developed. Culture revealed growths of *Penicillium* sp and *Acremonium* sp. Combined treatment with topical amphotericin B (0.1%) and natamycin, alternated hourly, and systemic ketoconazole resulted in resolution in a fluctuating course. The culture results were important for the diagnosis of a mixed infection and together with careful follow-up were critical for a good result.

Copyright © 2011, Buddhist Compassion Relief Tzu Chi Foundation. Published by Elsevier Taiwan LLC. All rights reserved.

1. Introduction

Fungal corneal ulcers or coinfection of bacteria and fungi are common in developing countries, among which *Aspergillus*, *Fusarium*, and *Candida* are the most common agents [1,2]. However, mixed fungal keratitis of two different genera has rarely been reported. Here, we reported a case of mixed fungal keratitis of *Penicillium* sp and *Acremonium* sp.

2. Case report

A 67-year-old male was injured in his right eye during the picking of herbs (*Ficus erecta* Thunb. var.). He came to our clinic for painful blurred vision 5 days later. On examination, biomicroscopy revealed no epithelial defect, but there was corneal stromal infiltration (two-thirds in depth) with a feathery border and an immunological reaction ring (Fig. 1A). A smear of corneal scraping showed the presence of hyphae (Fig. 1B). Topical natamycin (5%) hourly and oral voriconazole were prescribed immediately. However, hypopyon soon developed, which was resolved after

adding topical amphotericin B (0.1%), alternated hourly with natamycin, from Day 3 onward to the treatments. Oral voriconazole was changed into ketoconazole (400 mg/d) on Day 6 because of a skin rash and intolerable pruritus and increased hypopyon (Fig. 1C). Fungal culture with Sabouraud's dextrose agar showed growths of *Penicillium* species and *Acremonium* species (Fig. 1D and E). The hypopyon was completely resolved on Day 22. Amphotericin B was discontinued on Day 28 but was represcribed on Day 35 because of the reappearance of hypopyon. The lesion became almost a scar, and amphotericin B and ketoconazole was discontinued on Day 56. Unfortunately, a recurrence of an anterior chamber reaction was noted again on Day 58. Natamycin, amphotericin B, and systemic ketoconazole were used from this point for 2 more weeks until complete resolution (Fig. 1F). The final visual acuity was 20/40, and there was no recurrence in the following 6 months.

3. Discussion

Direct microscopy and culture results are both important tools in diagnosing infectious keratitis. Bharathi et al [3] even suggested a greater diagnostic value of KOH smear. However, diagnosis of a mixed infection depends on culture results. Mixed fungal keratitis has been rarely reported. Chin [4] reported a corneal perforation because of mixed infection of *Helminthosporium* sp and *Mima polymorpha*. In our patient, the corneal ulceration was caused by mixed infection of *Penicillium* sp and *Acremonium* sp. *Penicillium* and *Acremonium* sp are filamentous fungi widely spread in the

Conflict of interest: none.

* Corresponding author. Department of Ophthalmology, Buddhist Tzu Chi General Hospital, 707, Section 3, Chung-Yang Road, Hualien, Taiwan. Tel.: +886 3 856 1825x3253; fax: +886 3 857 8179.

E-mail address: yuanchieh.lee@gmail.com (Y.-C. Lee).




Fig. 1. (A) Slit lamp examination on referring to our clinic showed corneal stromal infiltration with a feathery border and an immunological reaction ring. (B) The smear of corneal scraping revealed septated fungal hyphae (white arrows). (C) Hypopyon increased suddenly on Day 6 of treatment with topical natamycin, amphotericin B, and oral voriconazole. (D and E) The culture of the corneal scraping demonstrated growths of *Penicillium* sp with brush-like phialides (D, black arrow heads) and *Acremonium* sp with conidia (E, dart arrows) aggregated at the apex of phialide. (F) Biomicroscopy showing complete resolution with faint scar.

environment, such as soil or tap water [5,6] and have been occasional etiologic pathogens of corneal ulceration, respectively [2]. *Acremonium* keratitis has been reported to be retractable and prone to recurrence and might require surgical interventions [7,8]. *Penicillium* keratitis has also led to penetrating keratoplasty [9]. The culture result of mixed infection of the above two fungi made us observe carefully for the corneal stromal infiltrate and ulceration, corneal edema, anterior chamber reaction, and also subjective symptoms.

Antifungal treatment is empirical conventionally [10]. It is not practical for the clinical ophthalmologist to ask a microbiology laboratory for identification and sensitivity studies to initiate and plan therapeutic management [11]. Many laboratories, including ours, are not equipped or experienced to perform sensitivity studies on fungi. Even laboratories with the necessary expertise may exhibit variability within their own results with the same fungi and may take days to weeks to perform the tests. Once obtained, the *in vitro* data and the clinical sensitivity response may not correlate [12]. It is most practical to consult published table of sensitivities for general groups of identified fungi against common available antifungal agents [12]. Careful clinical monitoring of the infiltrate and antifungal testing plays a role in the outcome of treatment [13]. Natamycin was reported the drug of choice for filamentous fungi keratitis, including *Acremonium* keratitis and *Penicillium* keratitis [2,14]. Natamycin has better penetration than amphotericin B in corneas with intact epithelium [15]. However, intensive topical natamycin application with oral voriconazole or ketoconazole was ineffective against the mixed *Acremonium* and *Penicillium* keratitis in our patient. The keratitis relapsed twice with an increased anterior chamber reaction after topical amphotericin B was discontinued. A regime involving a combination with topical amphotericin B seemed vital to cure. Therefore, we suggested that therapy for a fungal keratitis might be started with 5% natamycin hourly, but if the infection appeared to be progressing, then consideration should be given to adding or substituting another antifungal agent, such as amphotericin B or miconazole.

Fungal keratitis can easily become a surgical disease because of a delay in initiating medical treatment. Even with alertness and a prompt diagnosis of fungal infection, accurate culture results and a careful follow-up over the whole course are necessary for a good result.

References

- [1] Thomas PA. Fungal infections of the cornea. *Eye (Lond)* 2003;17:852–62.
- [2] Thomas PA. Current perspectives on ophthalmic mycoses. *Clin Microbiol Rev* 2003;16:730–97.
- [3] Bharathi MJ, Ramakrishnan R, Meenakshi R, Mittal S, Shivakumar C, Srinivasan M. Microbiological diagnosis of infective keratitis: comparative evaluation of direct microscopy and culture results. *Br J Ophthalmol* 2006;90:1271–6.
- [4] Chin GN. Corneal perforation due to *Helminthosporium* and *Mima* polymorpha. *Ann Ophthalmol* 1978;10:607–9.
- [5] Hedayati MT, Mohseni-Bandpi A, Moradi S. A survey on the pathogenic fungi in soil samples of potted plants from Sari hospitals, Iran. *J Hosp Infect* 2004;58:59–62.
- [6] Gonçalves AB, Paterson RR, Lima N. Survey and significance of filamentous fungi from tap water. *Int J Hyg Environ Health* 2006;209:257–64.
- [7] Wang MX, Shen DJ, Liu JC, Pflugfelder SC, Alfonso EC, Forster RK. Recurrent fungal keratitis and endophthalmitis. *Cornea* 2000;19:558–60.
- [8] Simonsz HJ. Keratomycosis caused by *Acremonium* Recifei, treated with keratoplasty, Miconazole and Ketoconazole. *Doc Ophthalmol* 1983;56:131–5.
- [9] Xie L, Dong X, Shi W. Treatment of fungal keratitis by penetrating keratoplasty. *Br J Ophthalmol* 2001;85:1070–4.
- [10] Loh AR, Hong K, Lee S, Mannis M, Acharya NR. Practice patterns in the management of fungal corneal ulcers. *Cornea* 2009;28:856–9.
- [11] Shadomy S. Susceptibility testing with antifungal drugs. In: Lennette EH, editor. *Manual of clinical microbiology*. 4th ed. Washington, DC: American Society for Microbiology; 1985. p. 991–9.
- [12] Alfonso EC, Forster RK, Garg P, Sharma S. Fungal infections. In: Foster CS, Azar DT, Cothman CH, editors. *Smolin and Thoft's the cornea: scientific foundations and clinical practice*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2005. p. 405–25.
- [13] Vemuganti GK, Garg P, Gopinathan U, Naduvilath TJ, John RK, Buddi R, et al. Evaluation of agent and host factors in progression of mycotic keratitis: a histologic and microbiologic study of 167 corneal buttons. *Ophthalmology* 2002;109:1538–46.
- [14] Kalavathy CM, Parmar P, Kaliyamurthy J, Philip VR, Ramalingam MD, Jesudasan CA, et al. Comparison of topical itraconazole 1% with topical natamycin 5% for the treatment of filamentous fungal keratitis. *Cornea* 2005;24:449–52.
- [15] O'Day DM, Head WS, Robinson RD, Clanton JA. Corneal penetration of topical amphotericin B and natamycin. *Curr Eye Res* 1986;5:877–82.